



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

VERY BRIGHT METEOR, MARCH 4, 1898.

OBSERVED BY H. D. CURTIS.

A very bright meteor was observed at College Park, March 4, 1898, 9<sup>h</sup> 50<sup>m</sup> 30<sup>s</sup> P. S. T., moving from  $\alpha = 13^{\text{h}} 40^{\text{m}}$ ,  $\delta = +25^{\circ}$  to  $\alpha = 15^{\text{h}} 40^{\text{m}}$ ,  $\delta = +40^{\circ}$ . Its path lay through the constellation *Bootes*, between the stars  $\beta$  and  $\delta$ . At a point just a little west of the line joining these two stars, there was a small but abrupt angle in its path, inclining towards the south. Several small meteors passed in almost exactly the same track during the next hour.

ELEMENTS AND EPHEMERIS OF COMET *b*, 1898  
(PERRINE).

BY R. T. CRAWFORD AND H. K. PALMER.

The following results were obtained from Mount Hamilton observations of March 20th and 22d and an observation taken at Berkeley on March 23d:—

$T = \text{March } 19.0580$ , G. M. T.

$i = 72^{\circ} 51' 42''$   
 $\Omega = 263 \ 15 \ 31$   
 $\omega = 49 \ 28 \ 52$  } Mean Equinox 1898.0.  
 $q = 1.1021$ .

(O.—C.)  $\Delta\lambda \cos \beta = -4''.7$   $\Delta\beta = +4''.7$ .

Constants to the equator:—

$x = [9.54097] \sin (27^{\circ} 37' 25'' + v) \sec^2 \frac{1}{2} v$ .  
 $y = [0.04218] \sin (295 \ 8 \ 42 + v) \sec^2 \frac{1}{2} v$ .  
 $z = [0.01954] \sin (24 \ 52 \ 17 + v) \sec^2 \frac{1}{2} v$ .

Ephemeris (Gr. Mean Midnight):—

1898.	APP. $\alpha$ .	APP. $\delta$ .	BRIGHTNESS.
March 29.5	21 <sup>h</sup> 55 <sup>m</sup> 9 <sup>s</sup>	+26° 29' 29"	0.99
April 2.5	22 12 4	30 26 49	0.96
6.5	22 29 56	34 12 41	0.91
10.5	22 48 45	37 43 53	0.86

The brightness is expressed in terms of the brightness at the time of discovery.

UNIVERSITY OF CALIFORNIA,  
STUDENTS' OBSERVATORY, March, 1898.